

# Nocturnal heart rate variability in obstructive sleep apnea syndrome: effect of automatic positive airway pressure

Iman H. Galal

**Background** The cyclic hypoxemia and arousal characteristic of obstructive sleep apnea (OSA) causes dysregulation in the autonomic nervous system (ANS) resulting in cardiac rhythm disturbances. Positive airway pressure therapy in OSA results in ANS regulation. This study aimed at investigating the effects of automatic positive airway pressure (APAP) over nocturnal heart rate variability (HRV) in OSA and to further identify this effect in hypertensive versus normotensive OSA patients.

**Patients and methods** A total of 25 patients (four with moderate and 21 with severe OSA) were included (male/female: 22/3; mean age 49.52±15.69 years). Patients were subdivided into two groups: 13 (52%) normotensive and 12 (48%) hypertensive patients. Two overnight sleep studies with polysomnographic-based nocturnal HRV were carried out: the first study was diagnostic, whereas the second one was an APAP titration study.

**Results** APAP results in significant reduction in most of time and frequency-domain indexes [except for standard deviation of average NN interval (SDANN,  $P=0.106$ ), HRV triangular index (HRV TI,  $P=0.057$ ), and high-frequency power (HF,  $P=0.109$ ); RR interval ( $P=0.024$ ), standard deviation of all RR intervals (SDNN,  $P=0.012$ ), mean of the standard deviation of all RR intervals for all 5-min segments (SDNN Index,  $P=0.009$ ), square root of the mean of the sum of the squares of differences between adjacent RR intervals

(RMSSD,  $P=0.002$ ), number of pairs of adjacent RR intervals differing by more than 50 ms in the entire analysis interval (NN50 count,  $P=0.004$ ), NN50 count divided by the total number of all RR intervals (%NN50,  $P=0.001$ ), total power ( $P=0.010$ ), very low-frequency power (VLF,  $P=0.018$ ), low-frequency power (LF,  $P=0.008$ ), and low frequency/high frequency (LF/HF,  $P=0.012$ ). HRV did not differ significantly between the two groups either before or after APAP; however, most of the time-domain and frequency-domain HRV indexes decreased significantly with APAP in the hypertensive group.

**Conclusion** Nocturnal HRV improved with APAP in OSA. APAP was more beneficial in terms of ANS regulation in OSA patients with coexisting hypertension than normotensive OSA patients.

*Egypt J Bronchol* 2017 11:372–378

© 2017 Egyptian Journal of Bronchology

*Egyptian Journal of Bronchology* 2017 11:372–378

**Keywords:** automatic positive airway pressure, frequency domain, heart rate variability, obstructive sleep apnea syndrome, time domain

Department of Chest Diseases, Ain Shams University, Cairo, Egypt

Correspondence to Iman H. Galal, MD, PhD, 4 Ibn Haggag Askalany Street, Heliopolis, Cairo - 11575, Egypt. Tel: +20 100 150 2179; fax: 21810643; e-mail: dr.imangalal@gmail.com

**Received** 23 April 2017 **Accepted** 11 May 2017

## Introduction

Positive airway pressure (PAP) is the standard treatment for patients with moderate-to-severe obstructive sleep apnea (OSA) – a sleep-related breathing disorder (SRBD) characterized by episodic complete or partial upper airway occlusion during sleep [1]. Over the past few years, the relationship between SRBD and cardiac arrhythmias have been thoroughly investigated [2–7]. Previous studies have shown that obstructive sleep apnea syndrome (OSAS) predisposes to clinically significant nocturnal impairment of the cardiac autonomic function, and such impairment correlated with the severity of OSAS [8–11]. The respiratory event-related cyclic hypoxemia and arousal characteristic of OSA causes dysregulation in the autonomic nervous system (ANS) in the form of enhanced sympathetic activity, and such impairment is thought to be the substrate necessary for heart rate variability (HRV) and consequently the development of cardiac arrhythmias [12–19].

HRV, the oscillation in the intervals separating consecutive heart beats, has gained importance as a noninvasive research and clinical tool for indirect quantitative and qualitative assessment of the cardiac

autonomic function in both health and disease [20]. HRV is mediated by three main mechanisms including vagal feedback from pulmonary stretch receptors, central medullary coupling respiratory and cardiovagal neurons, as well as the arterial baroreflex [21].

It is well known that OSAS is an independent risk factor for the development of systemic hypertension [22]. Altered autonomic control in the form of sympathetic activation represents a possible mechanism linking OSAS and hypertension [23].

Nasal continuous positive airways pressure therapy in OSA results in ANS regulation with subsequent improvement in HRV and nocturnal arrhythmias [24,25].

In view of the above, this study was undertaken in an attempt to investigate the effect of automatic positive

---

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work noncommercially, as long as the author is credited and the new creations are licensed under the identical terms.

airway pressure (APAP) on nocturnal HRV in patients with OSA and to further identify this effect in hypertensive compared with normotensive OSA patients.

## Patients and methods

### Study population

This prospective study included 25 randomly selected patients with polysomnographically (PSG) confirmed OSA. Participants were selected from among patients referred to the sleep laboratory with symptoms suggestive of SRBD. All included patients were newly diagnosed with OSAS and were never treated previously with PAP therapy. Patients were further subdivided into two groups based on their blood pressure values: the first group comprised 13 (52%) normotensive patients, whereas the second group comprised 12 (48%) hypertensive patients. All patients underwent two consecutive overnight sleep studies (PSG): the first study was diagnostic for OSA, whereas the second was a titration sleep study using APAP. Exclusion criteria were as follows: patients with atrial fibrillation, frequent ventricular premature beats, congestive heart failure, coronary artery disease, and stroke. Verbal consent was obtained from patients, and the study was approved by the Local Ethics Committee of the Faculty of Medicine at Ain Shams University.

### Polysomnography

Nocturnal, full-night PSG in the sleep laboratory starting from 10 p.m. to 6 a.m. was performed for all patients using a computerized system (N4000 Embla; Somnologica, Reykjavik, Iceland), including electroencephalogram, submental and anterior tibial electromyogram, oxygen saturation, ECG, inductance plethysmography of the chest and abdomen, nasal pressure sensor, and oronasal thermister monitoring. The recording was scored manually according to standard guidelines [26]. The apnea-hypopnea index (AHI) was calculated as the number of apneas and hypopneas per hour of total sleep time (TST). The threshold for diagnosis of OSA was set at an AHI of at least 5, and the severity of OSA was arbitrarily defined by the cutoff levels of AHI: of at least 5 to less than 15 episodes/h of TST for mild, of at least 15 to less than 30 episodes/h of TST for moderate, and of at least 30 episodes/h of TST for severe OSA [27].

### Automatic positive airway pressure titration study

The present study was performed during a standard, attended, overnight PSG in the sleep laboratory using an APAP device (AutoSet; ResMed, Sydney,

Australia) according to the recommendations of the American Academy of Sleep Medicine [28]. The AutoSet device is a computer-based PAP system that automatically modulates mask pressure in response to snoring, airflow limitation, or the presence of either apneas or hypopneas. The minimum pressure was set at 4 cmH<sub>2</sub>O, whereas the maximum pressure was set at 20 cmH<sub>2</sub>O according to the sleep medicine Task Force guidelines [29]. Only patients with successful APAP titration were included in the present study. The criteria for unsuccessful titration were as follows: TST of less than 4 h; TST in the supine position of less than 1 h; rapid eye movement (REM) sleep duration of less than 10 min [30]; excessive overshoots of delivered pressure associated with sleep disruption [31]; insufficient correction of obstructive events, not allowing us to recognize a therapeutic pressure; and median nocturnal air leak of more than 0.4 l/s [32,33].

### Heart rate variability analysis

Nocturnal ECG signals acquired by the PSG were digitalized. The analysis was carried out only for normal beats. HRV was evaluated using both time-domain and frequency-domain indexes. Time-domain indexes included the average length between each QRS complex (average RR interval), the standard deviation of all RR intervals (SDNN), the mean of the standard deviation of all RR intervals for all 5 min segments (SDNN index), the square root of the mean of the sum of the squares of differences between adjacent RR intervals (RMSSD), the number of pairs of adjacent RR intervals differing by more than 50 ms in the entire analysis interval (NN50 count), the NN50 count divided by the total number of all RR intervals [NN50 of total heart rate (HR) (%)], standard deviation of average NN interval (SDANN), and the total number of RR intervals divided by the maximum height of the histogram excluding boundaries (HRV triangular index). In the frequency-domain analysis, the power was calculated for very low-frequency (VLF, 0.0033–0.04 Hz), low-frequency (LF, 0.04–0.15 Hz), and high-frequency bands (HF, 0.15–0.4 Hz) as well as the LF/HF ratio [20].

### Statistical analysis

All statistical data analyses were carried out using the statistical package for social sciences software (SPSS for Windows, version 17.0; SPSS Inc, Chicago, Illinois, USA). Continuous numerical data are expressed as mean±SD, whereas categorical data are expressed as numbers and percentages. Normality distribution was tested using the Shapiro–Wilk test. Comparison between paired indexes was performed

using the paired *t*-test for parametric normally distributed data, and Wilcoxon's signed-rank test was used for nonparametric skewed data. Statistical significance was set at *P* values less than 0.05.

## Results

A total of 25 (22 males and three females) patients with OSAS with a mean age of 49.52±15.69 years and mean BMI of 39.14±8.00 kg/m<sup>2</sup> were included in this study; four patients had moderate OSAS, whereas the remaining 21 patients had severe OSAS. Patients with incomplete data were excluded. Table 1 shows the characteristics of all included patients as well as comparisons between different indexes before and during APAP. Several sleep indexes differed significantly during APAP: stages N1 and N2 nonrapid eye movement

sleep (NREM, *P*=0.000), stage N3 NREM sleep (*P*=0.000), REM stage (*P*=0.002), AHI (*P*=0.000), oxygen desaturation index (*P*=0.000), snoring index (*P*=0.000), arousal index (*P*=0.000), and REM latency (*P*=0.001). Both baseline and maximum HR decreased significantly with APAP (*P*=0.011 and 0.024, respectively). All time-domain indexes [except for SDANN (*P*=0.106) and HRV TI (*P*=0.057)] differed significantly during APAP: RR interval (*P*=0.024), SDNN (*P*=0.012), SDNN index (*P*=0.009), RMSSD (*P*=0.002), NN50 count (*P*=0.004), and %NN50 of total HR (*P*=0.001). All frequency-domain indexes [except for average HF power (*P*=0.109)] differed significantly during APAP: average total power (*P*=0.010), average VLF power (*P*=0.018), average LF power (*P*=0.008), and LF/HF ratio (*P*=0.012).

**Table 1 Patient characteristics**

Variables	OSAS (N=25)		<i>P</i>
	Before APAP	During APAP	
Age (years)	49.52±15.69		
Sex (male/female)	22/3		
BMI (kg/m <sup>2</sup> )	39.14±8.00		
TIB (min)	458.32±27.38	479.76±26.93	0.013
TST (min)	359.34±88.04	363.42±70.12	0.854
N1 and N2 (%)	77.54±13.60	51.56±14.49	0.000
N3 (%)	6.16±7.36	22.96±7.80	0.000
REM (%)	16.20±11.55	26.56±8.54	0.002
AHI (event/h of TST)	58.68±24.16	2.02±1.87	0.000
ODI (event/h of TST)	53.81±26.20	6.53±6.65	0.000
SI (% of TST)	41.66±22.42	10.23±9.74	0.000
Arl (number/h of TST)	50.68±21.28	11.57±6.15	0.000
SL (min)	20.72±17.25	25.52±33.25	0.686
RL (min)	178.19±91.45	89.38±68.41	0.001
Basal heart rate (beats/min)	77.60±11.98	74.32±12.65	0.011
Minimum heart rate (beats/min)	50.56±8.81	52.76±8.52	0.201
Maximum heart rate (beats/min)	94.92±10.05	90.28±11.44	0.024
RR interval (ms)	860.40±108.48	897.28±107.53	0.024
SDNN (ms)	113.84±56.80	95.12±51.66	0.012
SDNN index (ms)	90.52±61.03	69.16±55.12	0.009
RMSSD (ms)	86.88±84.55	69.28±84.87	0.002
NN50 count	5383.08±3822.95	3206.08±3352.36	0.004
NN50 of total HR (%)	19.39±15.40	11.07±12.06	0.001
SDANN (ms)	123.52±195.35	71.36±39.94	0.106
HRV TI	18.64±6.86	15.72±4.90	0.057
Average total power (ms <sup>2</sup> )	15792.35±12479.40	9757.16±3382.47	0.010
Average VLF power (ms <sup>2</sup> )	10098.24±10305.29	5834.48±3165.30	0.018
Average LF power (ms <sup>2</sup> )	3861.04±2865.53	2314.28±869.84	0.008
Average HF power (ms <sup>2</sup> )	1569.24±847.79	1389.96±988.58	0.109
LF/HF ratio	2.88±1.81	2.11±1.13	0.012

AHI, apnea-hypopnea index; APAP, automatic positive airway pressure; ArI, arousal index; HF, high frequency; HR, heart rate; HRV TI, heart rate variability triangular index; LF, low frequency; N1, stage 1 NREM sleep; N2, stage 2 NREM sleep; N3, stage 3 NREM sleep; NN50, the number of interval differences of successive NN intervals greater than 50 ms; NREM, nonrapid eye movement; ODI, oxygen desaturation index; OSAS, obstructive sleep apnea syndrome; REM, rapid eye movement sleep stage; RL, rapid eye movement latency; RMSSD, square root of the mean squared differences of successive NN intervals; SDANN, standard deviation of the average NN interval; SDNN, standard deviation of the NN interval; SI, snoring index; SL, sleep latency; VLF, very low frequency.

All time-domain and frequency-domain indexes were compared before and during APAP in both normotensive and hypertensive groups (Table 2). In the normotensive group, most of the time-domain and frequency-domain indexes did not differ significantly ( $P>0.05$ ) except for the RR interval ( $P=0.015$ ), average total power ( $P=0.028$ ), and average VLF power ( $P=0.013$ ). As for the hypertensive group, most of the time-domain and frequency-domain indexes differed significantly except for the RR interval ( $P=0.643$ ), SDANN ( $P=0.695$ ), and average VLF power ( $P=0.109$ ).

On comparing all time-domain and frequency-domain indexes before APAP in both groups, only SDNN

index differed significantly ( $P=0.04$ ), whereas all other time-domain and frequency-domain indexes did not differ significantly ( $P>0.05$ ). The same comparison during APAP showed no significant difference ( $P>0.05$ ) between the two groups with regard to all time-domain and frequency-domain indexes (Table 3).

## Discussion

Spectral analysis for nocturnal HRV in OSA patients before APAP use showed an increase in all frequency-domain indices, demonstrating both sympathetic and vagal activation. In addition, time-domain indexes suggestive of parasympathetic activation (namely

**Table 2 Heart rate variability before and during automatic positive airway pressure in normotensive and hypertensive obstructive sleep apnea syndrome patients**

Variables	Normotensive OSAS (N=13)			Hypertensive OSAS (N=12)		
	Before APAP	During APAP	P	Before APAP	During APAP	P
RR interval (ms)	825.08±111.88	887.62±75.93	0.015	898.67±94.59	907.75±136.77	0.643
SDNN (ms)	100.92±52.39	84.62±23.78	0.172	127.83±60.27	106.50±70.25	0.034
SDNN index (ms)	72.46±52.69	57.92±21.53	0.289	110.08±65.53	81.33±76.24	0.012
RMSSD (ms)	60.00±32.34	50.31±24.83	0.236	116.00±112.60	89.83±118.98	0.005
NN50 count	4708.92±3855.97	3624.15±3226.32	0.256	6113.42±3814.18	2753.17±3568.50	0.008
NN50 of total HR (%)	15.57±14.43	11.89±10.39	0.152	23.53±15.95	10.19±14.07	0.000
SDANN (ms)	88.46±40.17	65.38±30.05	0.056	161.50±280.14	77.83±49.07	0.695
HRV TI	17.77±8.55	16.38±4.81	0.603	19.58±4.60	15.00±5.12	0.002
Average total power (ms <sup>2</sup> )	17311.54±15835.91	10043.85±3725.33	0.028	14147.17±7780.26	9446.58±3101.66	0.026
Average VLF power (ms <sup>2</sup> )	11596.92±12409.25	5969.15±3056.24	0.013	8474.67±7628.13	5688.58±3409.62	0.109
Average LF power (ms <sup>2</sup> )	4188.62±3773.69	2562.54±1009.77	0.087	3506.17±1454.63	2045.33±621.89	0.005
Average HF power (ms <sup>2</sup> )	1352.08±550.50	1354.54±720.15	0.989	1804.50±1058.98	1428.33±1250.37	0.034
LF/HF ratio	3.23±2.08	2.22±1.24	0.071	2.51±1.46	1.99±1.03	0.043

APAP, automatic positive airway pressure; HF, high frequency; HR, heart rate; HRV TI, heart rate variability triangular index; LF, low frequency; NN50, the number of interval differences of successive NN intervals greater than 50 ms; OSAS, obstructive sleep apnea syndrome; RMSSD, square root of the mean squared differences of successive NN intervals; SDANN, standard deviation of average NN interval; SDNN, standard deviation of NN interval; VLF, very low frequency.

**Table 3 Heart rate variability in normotensive versus hypertensive obstructive sleep apnea syndrome patients before and during automatic positive airway pressure**

Variables	Before APAP (N=25)			During APAP (N=25)		
	Normotensive (n=13)	Hypertensive (n=12)	P	Normotensive (n=13)	Hypertensive (n=12)	P
RR interval (ms)	825.08±111.88	898.67±94.59	0.90	887.62±75.93	907.75±136.77	0.650
SDNN (ms)	100.92±52.39	127.83±60.27	0.077	84.62±23.78	106.50±70.25	0.852
SDNN index (ms)	72.46±52.69	110.08±65.53	0.04	57.92±21.53	81.33±76.24	0.574
RMSSD (ms)	60.00±32.34	116.00±112.60	0.152	50.31±24.83	89.83±118.98	0.574
NN50 count	4708.92±3855.97	6113.42±3814.18	0.320	3624.15±3226.32	2753.17±3568.50	0.406
NN50 of total HR (%)	15.57±14.43	23.53±15.95	0.168	11.89±10.39	10.19±14.07	0.347
SDANN (ms)	88.46±40.17	161.50±280.14	0.936	65.38±30.05	77.83±49.07	0.894
HRV TI	17.77±8.55	19.58±4.60	0.521	16.38±4.81	15.00±5.12	0.492
Average total power (ms <sup>2</sup> )	17311.54±15835.91	14147.17±7780.26	0.936	10043.85±3725.33	9446.58±3101.66	0.669
Average VLF power (ms <sup>2</sup> )	11596.92±12409.25	8474.67±7628.13	0.538	5969.15±3056.24	5688.58±3409.62	0.830
Average LF power (ms <sup>2</sup> )	4188.62±3773.69	3506.17±1454.63	0.852	2562.54±1009.77	2045.33±621.89	0.141
Average HF power (ms <sup>2</sup> )	1352.08±550.50	1804.50±1058.98	0.320	1354.54±720.15	1428.33±1250.37	0.611
LF/HF ratio	3.23±2.08	2.51±1.46	0.406	2.22±1.24	1.99±1.03	0.625

APAP, automatic positive airway pressure; HF, high frequency; HR, heart rate; HRV TI, heart rate variability triangular index; LF, low frequency; NN50, the number of interval differences of successive NN intervals greater than 50 ms; OSAS, obstructive sleep apnea syndrome; RMSSD, square root of the mean squared differences of successive NN intervals; SDANN, standard deviation of average NN interval; SDNN, standard deviation of NN interval; VLF, very low frequency.

RMSSD and %NN50 of total HR) were predominantly and significantly elevated before APAP, raising the assumption that OSAS can induce parasympathetic activation. Our findings of dual sympathetic and vagal activation in OSAS is not surprising, as other previous studies hypothesized that the repetitive upper airway collapse during sleep characteristic for OSAS causes initial vagal stimulation followed by abrupt sympathetic activation caused by hypoxemia and hypercapnia-related chemoreceptor stimulation [8,34].

On comparing all nocturnal HRV indexes before and during APAP, only SDANN and HRV TI time-domain indexes as well as HF power frequency-domain variables did not decrease significantly, suggesting that the reduction in sympathetic activity was more pronounced than that in vagal stimulation among OSAS patients in general, whether normotensive or hypertensive. A study by Karasulu *et al.* [35] using APAP reported that all time-domain indexes did not decrease significantly during APAP, whereas some frequency-domain indexes such as HF, LF, and VLF power decreased significantly on APAP, thus confirming the results of the present study regarding the reduction in both sympathetic and parasympathetic autonomic activities with APAP. The partial discrepancy between the results of both studies with regard to the time-domain indexes can be attributed to the difference in the type of APAP used in both studies; each APAP device has its own algorithm. The algorithm applied by the APAP device used in this study allows automatic increase or decrease in mask pressure in response to snoring, specific changes in inspiratory airflow contour morphology, or the presence of apnea or hypopnea in an attempt to restore complete airway patency. Patients with OSAS showed elevated nocturnal parasympathetic tone, as expressed by the higher values of %NN50 of total HR, RMSSD, and HF power, which reduced significantly with APAP, suggesting that APAP treatment in OSAS restores normal ANS activity. Chrysostomakis *et al.* [36] reported similar results following 2 months of treatment with CPAP. Moreover, a number of studies have also reported on the reduction in sympathetic activity in OSA patients on nasal continuous positive airways pressure [37–39].

A previous study conducted by the same author ([9]) on OSAS patients in an attempt to characterize the changes in nocturnal HRV and to further investigate the correlation between these changes and the severity of OSA results revealed clinically significant nocturnal impairment of the HRV mainly through spectral analysis and to a lesser extent through time-domain

indexes, and such impairment correlated with the severity of the disease.

The findings of the present study showed no significant differences in both time-domain and frequency-domain indices of HRV between the hypertensive and normotensive groups either before or during APAP. Although only the SDNN time-domain index was significantly higher before APAP among the hypertensive group compared with the normotensive group, it is noteworthy that in this study spectral analysis of HRV was considered as a more sensitive indicator of ANS function than the time-domain variables; this is simply because it is indicated to calculate the time-domain indexes over 24 h recordings, whereas spectral frequency-domain indexes can be calculated over a recording period of as short as 2–5 min [20]. Thus, based on the nature of the nocturnal recording in this study, the results of the frequency-domain indexes should be considered more seriously than those of the time-domain indexes. An interesting finding in this study was the elevated frequency-domain variables (except for HF power) in normotensive compared with hypertensive OSAS patients either before or during APAP titration, although not being significantly different. Thus, hypertensive OSAS patients exhibit increased parasympathetic activity (in terms of HF power) than normotensive OSAS patients. Cheng *et al.* [40] in his study demonstrated similar results with regard to nonsignificant differences in both time-domain and frequency-domain indexes of HRV between the two groups either before or during APAP treatment; however, their study reported lower HF power in hypertensive compared with normotensive OSAS patients. Accordingly, their study demonstrated decreased parasympathetic activity in hypertensive compared with normotensive OSAS patients either before or during APAP, and this was not in agreement with our results. A possible explanation for this discrepancy is the mismatch in BMI in both studies: OSA patients in the present study were far more obese, and it has been suggested that obesity affects autonomic activity.

When APAP was applied to normotensive patients, the frequency-domain indexes decreased but did not differ significantly. Although the VLF power increased significantly during APAP, the physiological explanation for the VLF index is not well established and seems to be related to the renin–angiotensin–aldosterone system, thermoregulation, and the peripheral vasomotor tone [41]. On the contrary, APAP application in hypertensive OSAS patients resulted in significant

changes in HRV: most of the HRV indexes decreased significantly with APAP. The increased sympathetic activity before APAP (as evidenced by increased LF power), the increased parasympathetic activity (as evidenced by increased HF power), as well as the increased sympathovagal activity of the heart (as evidenced by increased LF/HF ratio) significantly decreased with APAP, suggesting that hypertensive OSAS patients benefit from APAP treatment more than normotensive OSAS patients regarding ANS regulation. Moreover, it is worth mentioning that the elimination of airway collapse with APAP caused more significant reduction in LF compared with the HF and LF/HF ratio, suggesting that the degree of reduction in sympathetic activation with APAP was more than the reduction in vagal stimulation. Cheng *et al.* [40] and Heitmann *et al.* [42] revealed increased variability in autonomic activity during APAP titration in hypertensive versus normotensive OSAS patients. Moreover, the results of Cheng *et al.* [40] are in agreement with the present study regarding the predominance of LF changes of HRV among hypertensive patients during APAP titration.

There are several limitations to this study. First, only one APAP algorithm was investigated in this study. Second, HRV was investigated with one form of PAP therapy disregarding the effect of other forms of PAP therapy on HRV. Third, the immediate effect of APAP over HRV entails long-term follow-up to document such improvement with regular APAP usage.

It is worth mentioning that among the strengths in this study was the exclusion of OSAS patients with clinical conditions affecting autonomic activity and HRV. Moreover, the inclusion of normotensive OSAS patients for comparison with hypertensive patients clearly delineates the differences in HRV in normotensive versus hypertensive OSAS patients.

## Conclusion

Nocturnal HRV improved with APAP in OSA patients. Furthermore, APAP was more beneficial in terms of ANS regulation in OSA patients with coexisting hypertension than normotensive OSA patients.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

- 1 Kushida CA, Chediak A, Berry RB, Brown LK, Gozal D, Iber C, *et al.* Positive Airway Pressure Titration Task Force of the American Academy of Sleep Medicine. Clinical guidelines for the manual titration of positive airway pressure in patients with obstructive sleep apnea. *J Clin Sleep Med* 2008; **4**:157–171.
- 2 Tilkian AG, Guilleminault C, Schroeder JS, Lehrman KL, Simmons FB, Dement WC. Sleep-induced apnea syndrome: prevalence of cardiac arrhythmias and their reversal after tracheostomy. *Am J Med* 1977; **63**:348–358.
- 3 Zwillich C, Devlin T, White D, Douglas N, Weil J, Martin R. Bradycardia during sleep apnea: characteristics and mechanism. *J Clin Invest* 1982; **69**:1286–1292.
- 4 Guilleminault C, Connolly SJ, Winkle RA. Cardiac arrhythmia and conduction disturbances during sleep in 400 patients with sleep apnea syndrome. *Am J Cardiol* 1983; **52**:490–494.
- 5 Hoffstein V, Sateika S. Cardiac arrhythmias, snoring, and sleep apnea. *Chest* 1994; **106**:466–471.
- 6 Gami AS, Pressman G, Caples SM, Kanagala R, Gard JJ, Davison DE, *et al.* Association of atrial fibrillation and obstructive sleep apnea. *Circulation* 2004; **110**:364–367.
- 7 Mehra R, Benjamin EJ, Shahar E, Gottlieb DJ, Nawabit R, Kirchner HL, *et al.* Association of nocturnal arrhythmias with sleep-disordered breathing. *Am J Respir Crit Care Med* 2006; **173**:910–916.
- 8 Roche F, Gaspoz JM, Court-Fortune I, Minini P, Pichot V, Duverney D, *et al.* Screening of obstructive sleep apnea syndrome by heart rate variability. *Circulation* 1999; **100**:1411–1415.
- 9 Galal I. Nocturnal heart rate variability analysis as a screening tool for obstructive sleep apnea syndrome. *Egypt J Chest Dis Tuberc* 2012; **61**:187–195.
- 10 Park DH, Shin CJ, Hong SC, Yu J, Ryu S, Kim E, *et al.* Correlation between the severity of obstructive sleep apnea and heart rate variability indices. *J Korean Med Sci* 2008; **23**:226–231.
- 11 Aydin M, Altin R, Ozeren A, Kart L, Bilge M, Unalacak M. Cardiac autonomic activity in obstructive sleep apnea: time-dependent and spectral analysis of heart rate variability using 24-hour Holter electrocardiograms. *Tex Heart Inst J* 2004; **31**:132–136.
- 12 Wit AL, Rosen AR. After depolarizations and triggered activity. In: Fossard HA, Haber E, Jennings RB, Katz M, Morgan E, editors. *The heart and cardiovascular system*. New York, NY: Raven Press; 1986. pp.1449–1490.
- 13 Zipers DP, Inoue H. Autonomic neural control of cardiac excitable properties. In: Kulbertus HE, Franck G, editors. *Neurocardiology*. New York, NY: Futura; 1988. pp. 59–84.
- 14 Luenberger U, Jacob E, Sweer L, Waravdekar N, Zwillich C, Sinoway L. Surges of muscle sympathetic nerve activity during obstructive apnea are linked to hypoxemia. *J Appl Physiol* 1995; **79**:581–588.
- 15 Somers VK, Dyken ME, Clary MP, Abboud FM. Sympathetic neural mechanisms in obstructive sleep apnea. *J Clin Invest* 1995; **96**: 1897–1904.
- 16 Xie A, Skatrud JB, Puleo DS, Morgan BJ. Exposure to hypoxia produces long-lasting sympathetic activation in humans. *J Appl Physiol* 2001; **91**:1555–1562.
- 17 Narkiewicz K, van de Borne PJ, Pesek CA, Dyken ME, Montano N, Somers VK. Selective potentiation of peripheral chemoreflex sensitivity in obstructive sleep apnea. *Circulation* 1999; **99**: 1183–1189.
- 18 Dimsdale JE, Loreda JS, Profant J. Effect of continuous positive airway pressure on blood pressure: a placebo trial. *Hypertension* 2000; **35**:144–147.
- 19 Narkiewicz K, van de Borne PJ, Cooley RL, Dyken ME, Somers VK. Sympathetic activity in obese subjects with and without obstructive sleep apnea. *Circulation* 1998; **98**:772–776.
- 20 Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation* 1996; **93**:1043–1065.
- 21 Jo JA, Blasi A, Valladares E, Juarez R, Baydur A, Khoo MC. Determinants of heart rate variability in obstructive sleep apnea syndrome during wakefulness and sleep. *Am J Physiol Heart Circ Physiol* 2005; **288**:1103–1112.
- 22 Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med* 2000; **342**:1378–1384.

- 23 Fletcher EC. Sympathetic over activity in the etiology of hypertension of obstructive sleep apnea. *Sleep* 2003; **26**:15–19.
- 24 Liston R, Deegan P, McCreery C, McNicholas WT. Role of respiratory sleep disorders in the pathogenesis of nocturnal angina and arrhythmias. *Postgrad Med J* 1994; **70**:275–280.
- 25 Maser RE, Lenhard MJ, Rizzo AA, Vasile AA. Continuous positive airway pressure therapy improves cardiovascular autonomic function for persons with sleep-disordered breathing. *Chest* 2008; **133**:86–91.
- 26 Berry RB, Brooks R, Gamaldo CE, Harding SM, Loyd RM, Vaughn BV. *The AASM manual for the scoring of sleep and associated events: rules, terminology and technical specifications, version 2.3*. Darien, IL: American Academy of Sleep Medicine; 2016.
- 27 Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research: the Report of an American Academy of Sleep Medicine Task Force. *Sleep* 1999; **22**:667–689.
- 28 Littner M, Hirshkowitz M, Davila D, Anderson WM, Kushida CA, Woodson BT, *et al*. Standards of Practice Committee of the American Academy of Sleep Medicine. Practice parameters for the use of auto-titrating continuous positive airway pressure devices for titrating pressures and treating adult patients with obstructive sleep apnea syndrome. An American Academy of Sleep Medicine report. *Sleep* 2002; **25**:143–147.
- 29 Kushida CA, Chediak A, Berry RB, Brown LK, Gozal D, Iber C, *et al*. Clinical Guidelines for the Manual Titration of Positive Airway Pressure in Patients with Obstructive Sleep Apnea. Positive Airway Pressure Titration Task Force of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2008; **4**:157–171.
- 30 Marrone O, Insalaco G, Salvaggio A, Bonsignore G. Role of different nocturnal monitorings in the evaluation of CPAP titration by autoCPAP devices. *Respir Med* 2005; **99**:313–320.
- 31 Marrone O, Insalaco G, Bonsignore MR, Romano S, Salvaggio A, Bonsignore G. Sleep structure correlates of continuous positive airway pressure variations during application of an autotitrating continuous positive airway pressure machine in patients with obstructive sleep apnea syndrome. *Chest* 2002; **121**:759–767.
- 32 Teschler H, Berthon-Jones M. Intelligent CPAP systems, clinical experience. *Thorax* 1998; **53**:S49–S54.
- 33 Cross MD, Vennelle M, Engleman HM, White S, Mackay TW, Twaddle S, *et al*. Comparison of CPAP titration at home or the sleep laboratory in the sleep apnea hypopnea syndrome. *Sleep* 2006; **29**:1451–1455.
- 34 Narkiewicz K, Montano N, Cogliati C, van de Borne PJ, Dyken ME, Somers VK. Altered cardiovascular variability in obstructive sleep apnea. *Circulation* 1998; **98**:1071–1077.
- 35 Karasulu L, Epöztürk PO, Sökücü SN, Dalar L, Altin S. Improving Heart rate variability in sleep apnea patients: differences in treatment with auto-titrating positive airway pressure (APAP) versus conventional CPAP. *Lung* 2010; **188**:315–320.
- 36 Chrysostomakis SI, Simantirakis EN, Schiza SE, Karalis IK, Klapsinos NC, Siafakas NM, *et al*. Continuous positive airway pressure therapy lowers vagal tone in patients with obstructive sleep apnoea-hypopnoea syndrome. *Hellenic J Cardiol* 2006; **47**:13–20.
- 37 Jennum P, Wildschiodtz G, Christensen NJ, Schwartz T. Blood pressure, catecholamines, and pancreatic polypeptide in obstructive sleep apnea with and without nasal continuous positive airway pressure (nCPAP) treatment. *Am J Hypertens* 1989; **2**:847–852.
- 38 Minemura H, Akashiba T, Yamamoto H, Akahoshi T, Kosaka N, Horie T. Acute effects of nasal continuous positive airway pressure on 24-hour blood pressure and catecholamines in patients with obstructive sleep apnea. *Intern Med* 1998; **37**:1009–1013.
- 39 Hedner J, Darpo B, Eijnell H, Carlson J, Caidahl K. Reduction in sympathetic activity after long-term CPAP treatment in sleep apnoea: cardiovascular implications. *Eur Respir J* 1995; **8**:222–229.
- 40 Cheng JH, Hua CC, Chen NH, Liu YC, Yu CC. Autonomic activity difference during continuous positive airway pressure titration in patients with obstructive sleep apnea/hypopnea syndrome with or without hypertension. *Chang Gung Med J* 2011; **34**:410–417.
- 41 Vanderlei LC, Pastre CM, Hoshi RA, Carvalho TD, Godoy MF. Basic notions of heart rate variability and its clinical applicability. *Rev Bras Cir Cardiovasc* 2009; **24**:205–217.
- 42 Heitmann J, Ehlenz K, Penzel T, Becker HF, Grote L, Voigt KH, *et al*. Sympathetic activity is reduced by nCPAP in hypertensive obstructive sleep apnoea patients. *Eur Respir J* 2004; **23**:255–262.